

# A rule of thumb for cosmological backreaction

M. Parry

*Arnold Sommerfeld Center, Department für Physik,  
Ludwig-Maximilians-Universität München, Theresienstr. 37, D-80333 München, Germany*

(Dated: 10 May 2006)

In the context of second order perturbation theory, cosmological backreaction is seen to rescale both time and the scale factor. The issue of the homogeneous limit of long-wavelength perturbations is addressed and backreaction is quantified in terms of a gauge-invariant metric function that is the true physical degree of freedom in the homogeneous limit. The time integral of this metric function controls whether backreaction hastens or delays the expansion of the universe. As an example, late-time acceleration of the universe is shown to be inconsistent with a perturbative approach. Any tendency to accelerate the expansion requires negative non-adiabatic pressure fluctuations.

PACS numbers: 98.80.Cq

## I. INTRODUCTION

The application of cosmological perturbation theory to primordial fluctuations and the cosmic microwave background has been outstandingly successful [1]. Since inhomogeneities tend to grow in the universe, however, it is clear that first order perturbation theory will not remain valid. The question then arises: to what extent do these inhomogeneities act back on the expansion of the universe? Aligned with this question is the expectation that as observations become increasingly precise one might see second order effects at early times [2, 3, 4, 5, 6]. Consequently, the issue of cosmology perturbations to second order and beyond has been the subject of considerable inquiry [57].

The treatment of cosmological perturbations to second order was first carried out in the context of backreaction [7, 8]. This initiated the general study of inhomogeneities and averaging in cosmology [9, 10, 11, 12, 13], and culminated in a beautiful series of papers [14, 15, 16, 17] (see also [18]) highlighting the fact that there is both a kinematical backreaction because spatial averaging and time evolution do not commute, and a curvature backreaction because spatial averaging and spatial rescaling do not commute (see also [19]). At the very least, therefore, one should expect a rescaling of the cosmological parameters; in the extreme limit, the evolution of the universe can be mischaracterized by imposing on it the FLRW framework.

One of the deficiencies of the early work on second order perturbations was that it was not gauge-invariant and, hence, was difficult to interpret reliably. After the mathematical foundation of gauge-invariant perturbation theory to all orders was established [20, 21, 22], work began in earnest on quantifying backreaction and on identifying conservation laws for long-wavelength fluctuations [23, 24, 25, 26, 27, 28]. It was shown [22] that backreaction of long-wavelength modes behaves like a negative cosmological constant and it was anticipated that there could be a dynamical relaxation mechanism for the cosmological constant [29, 30]. Subsequently, there was some debate as to whether such backreaction was physi-

cally measurable [31, 32, 33, 34] and the analysis of backreaction was tightened. Specifically, it was noted [35] that one requires a physical clock, e.g. a secondary scalar field, to which a physical observable can be referred.

The most recent direction taken by the study of backreaction concerns the late-time acceleration of the universe [36, 37]. It suffices to say that arguments are forcefully put (e.g. [38, 39, 40, 41, 42, 43, 44, 45, 46]) both for and against the idea that backreaction can be responsible for late-time acceleration.

This paper is a conservative attempt to apply two lessons from the study of backreaction. The first is that the scale factor and other cosmological parameters deduced from observations are rescaled quantities and not the “bare” quantities entering the perturbative formalism. The second lesson is that if extremely long-wavelength modes are the primary contribution to backreaction, then it should be possible to understand their effect in terms of the local equations of homogeneous and isotropic cosmology [47]. Our approach is to suppose that a second order perturbative computation has been performed; the homogeneous component then represents the backreaction on the bare cosmological quantities.

In the next section, we deduce the general form of homogeneous scalar metric perturbations in an FLRW universe. We introduce the uniform curvature and conformal gauges in section III and construct the gauge invariant metric function that will be crucial in our discussion of backreaction. In section IV, we point out the weakness of the longitudinal gauge in the homogeneous limit and show that the uniform curvature gauge encodes the physical metric degree of freedom at all wavelengths. We also write down the perturbed Einstein equations for hydrodynamical matter in the uniform curvature gauge. In section V, treating homogeneous perturbations as arising at second order, we argue that backreaction is quantified by transforming to the conformal gauge and that, consequently, time and the scale factor are rescaled. We establish a rule of thumb for backreaction to hasten or delay the expansion of the universe, and illustrate what this implies for inflation and late-time acceleration.

## II. HOMOGENEOUS PERTURBATIONS

In the standard theory of cosmological perturbations [48, 49, 50, 51], all quantities are referred to the unperturbed manifold. Therefore, homogeneous perturbations are those with no dependence on the spatial coordinates of the background FLRW spacetime. One also assumes, usually implicitly, that the spatial average of first order perturbations vanishes. By this assumption, there can be no homogeneous perturbations at first order. For reasons that will become clear, we consider more general perturbations for which this assumption is relaxed. We will require only that the perturbations are bounded in space. For simplicity, we assume a flat model of the universe and use  $\eta$  to denote conformal time.

Homogeneous metric perturbations take the general form

$$\delta g_{\mu\nu} = a^2(\eta) \begin{bmatrix} 2\phi(\eta) & \Sigma_i(\eta) \\ \Sigma_j(\eta) & -H_{ij}(\eta) \end{bmatrix}, \quad (1)$$

where  $a(\eta)$  is the background scale factor. These can be decomposed into the usual scalar, vector and tensor pieces with respect to the spatial metric  $\delta_{ij}$  as follows [51]:

$$\begin{aligned} \Sigma_i &= S_i - B_{,i}, \\ H_{ij} &= -2\psi\delta_{ij} + 2E_{,ij} + F_{i,j} + F_{j,i} + h_{ij}, \end{aligned}$$

where  $S^i_{,i} = F^i_{,i} = h^{ij}_{,i} = h^i_{,i} = 0$ . The actual decomposition follows from the equations:

$$\begin{aligned} \nabla^2\psi &= 0, \\ \nabla^2 E &= 3\psi + \frac{1}{2}H^i_{,i}, \\ \nabla^2 B &= 0, \\ \nabla^2 F_j &= -4\psi_{,j}, \end{aligned}$$

where  $\nabla^2$  is the Laplacian operator on the 3-metric  $\delta_{ij}$ . The appropriate boundary conditions for these equations are that the quantities remain bounded. It then follows that all are simply functions of time and that the right hand sides must vanish. In particular, we have  $\psi = -\frac{1}{6}H^i_{,i}$ , and so  $S_i = \Sigma_i$  and  $h_{ij} = H_{ij} - \frac{1}{3}H^k_{,k}\delta_{ij}$ . We conclude that the scalar part of the metric perturbations has the form

$$\delta g_{\mu\nu}^{(S)} = a^2 \begin{bmatrix} 2\phi & 0 \\ 0 & 2\psi\delta_{ij} \end{bmatrix}. \quad (2)$$

## III. GAUGE INVARIANCE

From now on we will concentrate on scalar perturbations. As usual we will want to choose a gauge that exploits gauge-invariant quantities. It turns out that the longitudinal gauge is not a useful gauge here; instead we will consider the uniform curvature gauge and the conformal gauge.

Because  $B$  and  $E$  have no spatial dependence, they are effectively zero as far as the metric is concerned. But this does not mean we are in the longitudinal gauge. The coordinate transformation  $x^\mu \rightarrow \bar{x}^\mu = x^\mu + \delta x^\mu$  that affects the scalar perturbations has  $\delta x^\mu = (\xi^0, \xi^i)$  and leads to [51]

$$\begin{aligned} \phi &\rightarrow \bar{\phi} = \phi - \frac{1}{a}(a\xi^0)', \\ \psi &\rightarrow \bar{\psi} = \psi + \frac{a'}{a}\xi^0, \\ B &\rightarrow \bar{B} = B + \xi^0 - \xi', \\ E &\rightarrow \bar{E} = E - \xi, \end{aligned}$$

where  $'$  denotes a derivative with respect to  $\eta$ . We can choose the longitudinal gauge ( $\bar{E} = \bar{B} = 0$ ) in the usual way:  $\xi = E$ ,  $\xi^0 = E' - B$ , but this does not simplify the form of the metric at all. In fact, as long as  $\xi^0$  and  $\xi$  are functions of  $\eta$ ,  $B$  and  $E$  remain effectively zero. Since the choice of  $\xi$  has no bearing on  $\phi$  and  $\psi$ , we conclude that a residual degree of freedom, namely  $\xi^0$ , exists for homogeneous perturbations in the longitudinal gauge. Therefore, there will be a gauge artifact in our solutions if we simply let  $k \rightarrow 0$  in the standard treatment of cosmological perturbations. We will return to this point shortly.

A gauge-invariant quantity can be constructed solely out of  $\phi$  and  $\psi$ . The most convenient choice is

$$\Omega = \phi + \frac{1}{a} \left( \frac{a\psi}{\mathcal{H}} \right)', \quad (3)$$

where  $\mathcal{H} = a'/a$ . Furthermore, since perturbations of scalars transform as  $\delta q \rightarrow \bar{\delta q} = \delta q - q'_0 \xi^0$ ,

$$\delta q^{(gi)} = \delta q + \frac{q'_0 \psi}{\mathcal{H}} \quad (4)$$

will also be gauge-invariant. It is straightforward to show that in terms of the gauge-invariant quantities natural to the longitudinal gauge  $\delta q^{(gi)} = \delta q^{(LG)} + q'_0 \Psi / \mathcal{H}$ .

### A. Uniform curvature gauge

Choosing  $\xi^0 = -\psi/\mathcal{H}$ , completely fixes the gauge and gives  $\bar{\psi} = 0$  and  $\bar{\phi} = \Omega$ . Therefore, only the  $g_{00}$  component of the metric is perturbed and all scalar perturbations are automatically gauge-invariant. Because  $\bar{\psi} = 0$  this gauge is called the uniform curvature gauge or the spatially flat gauge.

A perfect fluid is characterized by its density  $\rho$ , pressure  $p$  and 3-velocity  $u^i$ . In the uniform curvature gauge and to first order in homogeneous perturbations, the Einstein equations become

$$-3\mathcal{H}^2\Omega = 4\pi G a^2 \delta\rho^{(gi)}, \quad (5)$$

$$0 = \delta u_i^{(gi)}, \quad (6)$$

$$(2\mathcal{H}' + \mathcal{H}^2)\Omega + \mathcal{H}\Omega' = 4\pi G a^2 \delta p^{(gi)}, \quad (7)$$

where  $\delta u_i^{(gi)} = \delta u_i + a\psi_{,i}/\mathcal{H} = \delta u_i^{(LG)} + a\Psi_{,i}/\mathcal{H}$ . Making use of the background equations  $3\mathcal{H}^2 = 8\pi G a^2 \rho_0$ ,  $\mathcal{H}' - \mathcal{H}^2 = -4\pi G a^2(\rho_0 + p_0)$  and  $\rho_0' + 3\mathcal{H}(\rho_0 + p_0) = 0$ , we can rewrite the first and third equations as

$$\Omega = -\frac{1}{2}\frac{\delta\rho^{(gi)}}{\rho_0}, \quad (8)$$

$$\delta\rho^{(gi)'} = -3\mathcal{H}(\delta\rho^{(gi)} + \delta p^{(gi)}). \quad (9)$$

As an example, we consider the case of adiabatic perturbations:  $\delta p^{(gi)} = (p_0'/\rho_0')\delta\rho^{(gi)}$ . Then

$$\Omega = -\frac{3}{2}\zeta\left(1 + \frac{p_0}{\rho_0}\right), \quad (10)$$

where

$$\zeta \equiv -\frac{\mathcal{H}\delta\rho^{(gi)}}{\rho_0'} = -\Psi - \frac{\mathcal{H}\delta\rho^{(LG)}}{\rho_0'} \quad (11)$$

is the well known invariant from cosmological perturbation theory. Here  $\zeta = \text{const.}$

### B. Conformal gauge

If we choose  $\xi^0 = \int d\eta(\phi + \psi)$ , then  $\bar{\phi} = -\bar{\psi}$  and the metric is returned to a conformal FLRW metric. Although the conformal gauge is not a good gauge— $\xi^0$  is only known up to an additive constant—it suggests a way to quantify the backreaction of perturbations on the background scale factor. We will develop the idea fully in section V.

## IV. A COMMENT ON $k = 0$ VS. $k \rightarrow 0$

The standard treatment of (inhomogeneous) scalar perturbations shows there is essentially one metric scalar degree of freedom: there are two gauge-invariant scalars with one constraint. To be precise, there is a combination of the two scalars, call it  $D$ , which satisfies a set of purely spatial differential equations. The  $k \rightarrow 0$  prescription is to find  $D$  and then set the spatial derivatives to zero in the remaining equations. On the other hand, choosing  $k = 0$  means dropping all spatial derivatives from the start and so there is no hope of subsequently finding  $D$ . The weakness of the longitudinal gauge is that it requires knowledge of  $D$  to be able to fully specify homogeneous perturbations. This is exactly the gauge artifact mentioned previously. Fortunately, as we now show, the uniform curvature gauge is independent of  $D$  in the homogeneous limit. In other words,  $k = 0$  and  $k \rightarrow 0$  are the same in the uniform curvature gauge. We will illustrate this fact by again considering the case of a perfect fluid.

The uniform curvature gauge can be extended to general perturbations [49, 50, 52, 53] by choosing  $\xi = E$  so that  $\bar{E} = 0$  in addition to  $\bar{\psi} = 0$ . In this way, the scalar

metric perturbations are confined solely to the time-time and time-space components of the metric. We also introduce the gauge-invariant combination

$$\beta = B - E' - \frac{\psi}{\mathcal{H}}. \quad (12)$$

To facilitate comparison with the longitudinal gauge, in which the scalar perturbations  $\phi$  and  $\psi$  become the usual gauge-invariant quantities  $\Phi$  and  $\Psi$ , we note that

$$\left. \begin{aligned} \Phi &= \Omega + \beta' + \mathcal{H}\beta \\ \Psi &= -\mathcal{H}\beta \end{aligned} \right\} \iff \left\{ \begin{aligned} \Omega &= \Phi + \Psi + \left(\frac{\Psi}{\mathcal{H}}\right)' \\ \beta &= -\frac{\Psi}{\mathcal{H}} \end{aligned} \right. \quad (13)$$

The perturbed equations for a perfect fluid in the uniform curvature gauge are quite compact:

$$-3\mathcal{H}^2\Omega - \mathcal{H}\nabla^2\beta = 4\pi G a^2 \delta\rho^{(gi)}, \quad (14)$$

$$\mathcal{H}\Omega_{,i} = 4\pi G a(\rho_0 + p_0)\delta u_i^{(gi)}, \quad (15)$$

$$\left[(2\mathcal{H}' + \mathcal{H}^2)\Omega + \mathcal{H}\Omega' + \frac{1}{2}\nabla^2 D\right]\delta_j^i - \frac{1}{2}D^{,i}_{,j} = 4\pi G a^2 \delta p^{(gi)}\delta_j^i, \quad (16)$$

where  $D = \Omega + \beta' + 2\mathcal{H}\beta = \Phi - \Psi$ . The off-diagonal terms and the differences between the diagonal terms of (16) imply that, for  $D$  bounded,  $D = D(\eta)$ . If the spatial average of  $D$  is to vanish then we must have  $D = 0$ . However, in the homogeneous limit of the uniform curvature gauge, we do not need to know  $D$ . This is because the term in  $\beta$  also drops out; we return to eqns. (5)-(7) and  $\Omega$  is the one metric degree of freedom required. In contrast, in the longitudinal gauge, terms in both  $\Phi$  and  $\Psi$  remain if  $k = 0$ . We conclude that  $\Omega$  is the true physical degree of freedom describing long-wavelength fluctuations and that the discussion of homogeneous perturbations is best had in the uniform curvature gauge. The uniform curvature gauge can also be extended to second order [3, 24, 26, 28].

Pertinent to this discussion, one of the conservation laws that can be derived from the above equations is

$$\zeta' - \frac{1}{3}\nabla^2\beta - \frac{1}{3}a^{-1}\nabla^i\delta u_i^{(gi)} = -\frac{\mathcal{H}}{\rho_0 + p_0}\delta p_{\text{nad}}, \quad (17)$$

where  $\delta p_{\text{nad}} = \delta p^{(gi)} - (p_0'/\rho_0')\delta\rho^{(gi)}$  is the non-adiabatic pressure fluctuation.

## V. COSMOLOGICAL BACKREACTION

All cosmological observations are carried out assuming an underlying FLRW metric. The logic of backreaction is that, while perturbation theory remains valid, higher order quantities affect the evolution of lower order quantities. As emphasized earlier, this implies, for example, that  $a(\eta)$  is the bare scale factor; the actual observed scale factor is found after backreaction has been taken into account. We are looking for the homogeneous effect of second order terms on background quantities.

In this light, the homogeneous perturbations in section II should be thought of as second order perturbations; first order quantities vanish in the homogeneous limit. As an example, if  $q = q_0 + q_1 + q_2 + \dots$  is a perturbative expansion then, generically, we would expect the homogeneous perturbation

$$\delta q \sim \langle q_1^2 \rangle.$$

Note that this does not vanish, in contrast to the term  $\langle q_0 q_2 \rangle = q_0 \langle q_2 \rangle = 0$ . Furthermore, the observed scale factor can then be found by converting to the conformal gauge. Unfortunately, because the conformal gauge is not fixed, this procedure is not unique. We claim that it is most consistent to perform computations in the uniform curvature gauge and from there convert to the conformal gauge ( $\phi \rightarrow \Omega$ ,  $\psi \rightarrow 0$ ).

To be specific,  $\xi^0 = \int d\eta \Omega$  and all background quantities should be written in terms of the rescaled conformal time  $\bar{\eta}$  via

$$\eta = \bar{\eta} - \int d\eta \Omega(\eta) \simeq \bar{\eta} - \int d\bar{\eta} \Omega(\bar{\eta}). \quad (18)$$

Such rescaling of the time is a common technique in nonlinear perturbation theory and is similar to the approaches found in [5, 47, 54]. The fact that  $\xi^0$  is only specified up to a constant is not a serious drawback; it will be fixed once the initial time is chosen, e.g. when  $a(\eta_0) = 1$ . Apart from the rescaling of the time, this integration constant does not affect the scale factor nor the Hubble and deceleration parameters:

$$\bar{a}(\bar{\eta}) = a(\eta) = a(\bar{\eta} - \xi^0), \quad (19)$$

$$\bar{\mathcal{H}}(\bar{\eta}) = \mathcal{H}(\eta) (1 - \Omega), \quad (20)$$

$$\bar{q}(\bar{\eta}) = q(\eta) + \frac{\Omega'}{\mathcal{H}(\eta)}. \quad (21)$$

Equation (18) suggests a useful rule of thumb: cosmological evolution will be delayed (hastened) when  $\int d\eta \Omega(\eta)$  is positive (negative).

To add some physical content to this statement, let us suppose second order effects can be treated like a perfect fluid component. Such a situation was shown to arise in inflationary backreaction [22, 34, 35]: the second order terms contribute to an effective energy-stress-momentum tensor that is added to the right hand side of the unperturbed Einstein equations. In this particular case,  $\delta p^{(gi)} \simeq -\delta \rho^{(gi)} \simeq \text{const.} > 0$ . Since  $\rho_0$  is also roughly constant but positive during inflation, we have from eq. (5),

$$\int d\eta \Omega \simeq -\frac{1}{2} \frac{\delta \rho^{(gi)}}{\rho_0} \eta > 0. \quad (22)$$

We conclude, as in [22], that backreaction works to slow inflationary expansion.

A question currently of some interest is whether backreaction can lead to late-time acceleration of the universe. Since a stage of matter domination has  $q = \frac{1}{2}$  and de Sitter expansion  $q = -1$ , we can already see that such a transition would necessarily require the breakdown of perturbation theory. Nevertheless, our prescription can indicate the tendency of backreaction to accelerate the expansion of the universe. Taking  $p_0 = 0$  for dust so that the background scale factor is  $a \sim \eta^2$ , we can say for certain that we require negative non-adiabatic pressure fluctuations (see also [39, 54]):

$$\bar{q} \simeq \frac{1}{2} + \frac{\Omega'}{\mathcal{H}} = \frac{1}{2} + \frac{3}{2} \frac{\delta p_{\text{nad}}}{\rho_0}. \quad (23)$$

This should also be combined with  $\delta \rho^{(gi)} > 0$  so that  $\int d\eta \Omega < 0$  is guaranteed.

## VI. CONCLUSIONS

Since first order cosmological fluctuations are required to vanish under spatial averaging, homogeneous perturbations can only arise at second order. They then represent corrections to the background quantities. We formulated a gauge-invariant description of homogeneous perturbations in order to quantify this backreaction. The uniform curvature gauge is the most convenient gauge to exploit this gauge invariance and we found that the gauge-variant metric function  $\Omega$  is the true physical degree of freedom in the homogeneous limit. The final step is to transform to the conformal gauge to deduce the actual observed background quantities. We saw that this amounted to a rescaling of the time coordinate, an outcome which is well known in nonlinear perturbation theory. The value of the rule of thumb presented here is that once second order perturbations have been computed—admittedly a non-trivial task—one can quickly see what effect they have on the evolution of the universe. The gauge invariant quantity  $\Omega$  also clarifies issues surrounding  $k = 0$  vs.  $k \rightarrow 0$  and suggests new metric ansätze to study fully nonlinear perturbations.

## Acknowledgments

It is a pleasure to thank L. R. W. Abramo, T. Buchert and S. Winitzki for their insightful comments. MP was supported by Sonderforschungsbereich SFB 375 für Astro-Teilchenphysik der Deutschen Forschungsgemeinschaft.

- Nucl. Phys. **B667**, 119 (2003), astro-ph/0209156.
- [3] N. Bartolo, E. Komatsu, S. Matarrese, and A. Riotto, Phys. Rept. **402**, 103 (2004), astro-ph/0406398.
  - [4] P. Martineau and R. H. Brandenberger, Phys. Rev. **D72**, 023507 (2005), astro-ph/0505236.
  - [5] N. Bartolo, S. Matarrese, and A. Riotto, JCAP **0508**, 010 (2005), astro-ph/0506410.
  - [6] F. Finelli, G. Marozzi, G. P. Vacca, and G. Venturi (2006), gr-qc/0604081.
  - [7] T. Futamase, Phys. Rev. Lett. **61**, 2175 (1988).
  - [8] T. Futamase, Mon. Not. R. astr. Soc. **237**, 187 (1989).
  - [9] S. Bildhauer and T. Futamase, Gen. Rel. Grav. **23**, 1251 (1991).
  - [10] R. M. Zalaletdinov, Gen. Rel. Grav. **24**, 1015 (1992).
  - [11] T. Buchert and J. Ehlers, Astron. Astrophys. **320**, 1 (1997), astro-ph/9510056.
  - [12] H. Russ, M. H. Soffel, M. Kasai, and G. Börner, Phys. Rev. **D56**, 2044 (1997), astro-ph/9612218.
  - [13] R. M. Zalaletdinov, Bull. Astron. Soc. India **25**, 401 (1997), gr-qc/9703016.
  - [14] T. Buchert, Gen. Rel. Grav. **32**, 105 (2000), gr-qc/9906015.
  - [15] T. Buchert, Gen. Rel. Grav. **33**, 1381 (2001), gr-qc/0102049.
  - [16] T. Buchert and M. Carfora, Class. Quant. Grav. **19**, 6109 (2002), gr-qc/0210037.
  - [17] T. Buchert and M. Carfora, Phys. Rev. Lett. **90**, 031101 (2003), gr-qc/0210045.
  - [18] G. F. R. Ellis and T. Buchert, Phys. Lett. **A347**, 38 (2005), gr-qc/0506106.
  - [19] M. Mars and R. M. Zalaletdinov, J. Math. Phys. **38**, 4741 (1997), dg-ga/9703002.
  - [20] V. F. Mukhanov, L. R. W. Abramo, and R. H. Brandenberger, Phys. Rev. Lett. **78**, 1624 (1997), gr-qc/9609026.
  - [21] M. Bruni, S. Matarrese, S. Mollerach, and S. Sonego, Class. Quant. Grav. **14**, 2585 (1997), gr-qc/9609040.
  - [22] L. R. W. Abramo, R. H. Brandenberger, and V. F. Mukhanov, Phys. Rev. **D56**, 3248 (1997), gr-qc/9704037.
  - [23] G. Rigopoulos, Class. Quant. Grav. **21**, 1737 (2004), astro-ph/0212141.
  - [24] D. H. Lyth and D. Wands, Phys. Rev. **D68**, 103515 (2003), astro-ph/0306498.
  - [25] H. Noh and J.-c. Hwang, Phys. Rev. **D69**, 104011 (2004).
  - [26] K. A. Malik and D. Wands, Class. Quant. Grav. **21**, L65 (2004), astro-ph/0307055.
  - [27] F. Vernizzi, Phys. Rev. **D71**, 061301 (2005), astro-ph/0411463.
  - [28] K. A. Malik, JCAP **0511**, 005 (2005), astro-ph/0506532.
  - [29] R. H. Brandenberger (1999), hep-th/0004016.
  - [30] R. H. Brandenberger (2002), hep-th/0210165.
  - [31] W. Unruh (1998), astro-ph/9802323.
  - [32] L. R. Abramo and R. P. Woodard, Phys. Rev. **D65**, 063515 (2002), astro-ph/0109272.
  - [33] N. Afshordi and R. H. Brandenberger, Phys. Rev. **D63**, 123505 (2001), gr-qc/0011075.
  - [34] G. Geshnizjani and R. Brandenberger, Phys. Rev. **D66**, 123507 (2002), gr-qc/0204074.
  - [35] G. Geshnizjani and R. Brandenberger, JCAP **0504**, 006 (2005), hep-th/0310265.
  - [36] A. G. Riess et al. (Supernova Search Team), Astron. J. **116**, 1009 (1998), astro-ph/9805201.
  - [37] S. Perlmutter et al. (Supernova Cosmology Project), Astrophys. J. **517**, 565 (1999), astro-ph/9812133.
  - [38] E. W. Kolb, S. Matarrese, A. Notari, and A. Riotto, Phys. Rev. **D71**, 023524 (2005), hep-ph/0409038.
  - [39] E. W. Kolb, S. Matarrese, A. Notari, and A. Riotto, Mod. Phys. Lett. **A20**, 2705 (2005), astro-ph/0410541.
  - [40] E. E. Flanagan, Phys. Rev. **D71**, 103521 (2005), hep-th/0503202.
  - [41] G. Geshnizjani, D. J. H. Chung, and N. Afshordi, Phys. Rev. **D72**, 023517 (2005), astro-ph/0503553.
  - [42] A. Ishibashi and R. M. Wald, Class. Quant. Grav. **23**, 235 (2006), gr-qc/0509108.
  - [43] T. Buchert, Class. Quant. Grav. **23**, 817 (2006), gr-qc/0509124.
  - [44] S. Räsänen, Class. Quant. Grav. **23**, 1823 (2006), astro-ph/0504005.
  - [45] E. R. Siegel and J. N. Fry, Astrophys. J. **628**, L1 (2005), astro-ph/0504421.
  - [46] P. Martineau and R. Brandenberger (2005), astro-ph/0510523.
  - [47] R. H. Brandenberger and C. S. Lam (2004), hep-th/0407048.
  - [48] E. M. Lifshitz and I. M. Khalatnikov, Adv. Phys. **12**, 185 (1963).
  - [49] J. M. Bardeen, Phys. Rev. **D22**, 1882 (1980).
  - [50] H. Kodama and M. Sasaki, Prog. Theor. Phys. Suppl. **78**, 1 (1984).
  - [51] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, Phys. Rept. **215**, 203 (1992).
  - [52] V. N. Lukash, Sov. Phys. JETP **52**, 807 (1980).
  - [53] D. H. Lyth, Phys. Rev. **D31**, 1792 (1985).
  - [54] Y. Nambu, Phys. Rev. **D71**, 084016 (2005), gr-qc/0503111.
  - [55] R. Easther and M. Parry, Phys. Rev. **D62**, 103503 (2000), hep-ph/9910441.
  - [56] D. Langlois and F. Vernizzi, Phys. Rev. **D72**, 103501 (2005), astro-ph/0509078.
  - [57] Space does not allow us to mention the work on fully non-linear cosmological perturbations. The interested reader may turn to [33, 55, 56] and references therein.